arXiv:math/0703847v2 [math.SP] 17 Feb 2008

The relation between the counting function $N(\lambda)$ and the heat kernel $K(t)$

BY WU-SHENG DAI and MI XIE *

Abstract

For a given spectrum $\{\lambda_n\}$ of the Laplace operator on a Riemannian manifold, in this paper, we present a relation between the counting function $N(\lambda)$, the number of eigenvalues (with multiplicity) smaller than λ, and the heat kernel $K(t)$, defined by $K(t) = \sum_n e^{-\lambda_n t}$. Moreover, we also give an asymptotic formula for $N(\lambda)$ and discuss when $\lambda \to \infty$ in what cases $N(\lambda) = K(1/\lambda)$.

The relation between the spectrum of the Laplace operator on a Riemannian manifold and the geometry of this Riemannian manifold is an important subject [1, 2, 3, 4], and the problem of spectral asymptotics is one of the central problems in the theory of partial differential operators [5]. For a given spectrum $\{\lambda_n\}$ of the Laplace operator on a Riemannian manifold, one can in principle obtain the counting function $N(\lambda)$, defined to be

$$N(\lambda) = \text{the number of eigenvalues (with multiplicity) of the Laplace operator smaller than } \lambda, \tag{1}$$

and the heat kernel, defined to be

$$K(t) = \sum_n e^{-\lambda_n t}. \tag{2}$$

One of the main problems is to seek the asymptotic expansions of $N(\lambda)$ and $K(t)$. Usually, it is relatively easy to obtain the asymptotic expansion of the heat kernel $K(t)$. Nevertheless, it is difficult to calculate the asymptotic expansion of the counting function $N(\lambda)$ [2]. The Hardy-Littlewood-Karamata Tauberian theorem gives the first term of the asymptotic expansion of $N(\lambda)$ [6], but does not provide any information beyond the first-order term. In this paper, we point out a relation between $N(\lambda)$ and $K(t)$.

---

*We are very indebted to Dr G. Zeitrauman for his encouragement. This work is supported in part by NSF of China, under Project No.10605013 and No.10675088.

Theorem 1

$$K\left(t\right)=t\int_{0}^{\infty}N\left(\lambda\right)e^{-\lambda t}d\lambda. \tag{3}$$

Proof. The generalized Abel partial summation formula reads

$$\sum_{u_1<\lambda_n\leq u_2}b\left(n\right)f\left(\lambda_n\right)=B\left(u_2\right)f\left(u_2\right)-B\left(u_1\right)f\left(u_1\right)-\int_{u_1}^{u_2}B\left(u\right)f'\left(u\right)du, \tag{4}$$

where $\lambda_i\in\mathbb{R}$, $\lambda_1\leq\lambda_2\leq\cdots\leq\lambda_n\leq\cdots$, and $\lim_{n\to\infty}\lambda_n=\infty$. $f\left(u\right)$ is a continuously differentiable function on $[u_1,u_2]$ $(0\leq u_1<u_2,\ \lambda_1\leq u_2)$, $b\left(n\right)$ $(n=1,2,3,\cdots)$ are arbitrary complex numbers, and $B\left(u\right)=\sum_{\lambda_n\leq u}b\left(n\right)$. We apply the generalized Abel partial summation formula, Eq. (4), with $f\left(u\right)=e^{-u(s-s_0)}$ and $b\left(n\right)=a_ne^{-\lambda_ns_0}$, where s, $s_0\in\mathbb{C}$. Then

$$A\left(u_2,s\right)-A\left(u_1,s\right)=A\left(u_2,s_0\right)e^{-u_2(s-s_0)}-A\left(u_1,s_0\right)e^{-u_1(s-s_0)}+\left(s-s_0\right)\int_{u_1}^{u_2}A\left(u,s_0\right)e^{-u(s-s_0)}du, \tag{5}$$

where

$$A\left(u,s\right)=\sum_{\lambda_n\leq u}a_ne^{-\lambda_ns}. \tag{6}$$

Setting $a_n=1$ in Eq. (6), we find

$$A\left(\lambda,0\right)=\sum_{\lambda_n\leq\lambda}1=N\left(\lambda\right),$$

the counting function, and

$$A\left(\infty,t\right)=\sum_{n}e^{-\lambda_nt}=K\left(t\right),$$

the heat kernel. By Eq. (6), we also have $A\left(0,t\right)=0$. Then, by Eq. (5), we have

$$K\left(t\right)=A\left(\infty,t\right)-A\left(0,t\right)=t\int_{0}^{\infty}N\left(\lambda\right)e^{-\lambda t}d\lambda. \tag{7}$$

This is just Eq. (3). ■

Furthermore we can also obtain the following theorem.

Theorem 2

$$N\left(\lambda\right)=\frac{1}{2\pi i}\int_{c-i\infty}^{c+i\infty}K\left(t\right)\frac{e^{\lambda t}}{t}dt,\qquad c>\lim_{n\to\infty}\frac{\ln n}{\lambda_n}. \tag{8}$$

Proof. By the Perron formula, we have

$$\sum_{\mu_n<x}a_n=\frac{1}{2\pi i}\int_{c-i\infty}^{c+i\infty}f\left(t\right)\frac{x^t}{t}dt, \tag{9}$$

where

$$f\left(s\right)=\sum_{n=1}^{\infty}\frac{a_n}{\mu_n^s}, \tag{10}$$

and c is a constant which is greater than the abscissa of absolute convergence of the Dirichlet series $f(s)$. Setting

$$a_n = 1 \text{ and } \mu_n = e^{\lambda_n}$$

in Eq. (10), we obtain the heat kernel,

$$f(t) = \sum_{n=1}^{\infty} e^{-\lambda_n t} = K(t).$$

The abscissa of absolute convergence of $f(t)$ equals its abscissa of convergence, equaling $\overline{\lim}_{n\to\infty} \ln n/\lambda_n = \lim_{n\to\infty} \ln n/\lambda_n$. Thus, by Eq. (9), we have

$$N(\lambda) = \sum_{\lambda_n<\lambda} 1 = \frac{1}{2\pi i}\int_{c-i\infty}^{c+i\infty} K(t)\frac{e^{\lambda t}}{t}dt,$$

and $c > \lim_{n\to\infty} \ln n/\lambda_n$. This proves the theorem. ■

The above two theorems give the relation between the counting function $N(\lambda)$ and the heat kernel $K(t)$.

One of the reasons why the counting function $N(\lambda) = \sum_{\lambda_n<\lambda} 1$ is very difficult to calculate is that one often encounters some unsolved problems in number theory when calculating $N(\lambda)$. For example, when calculating the counting function for the spectrum of the Laplace operator on a tori, one encounters the Gauss circle problem in number theory [2]. In the following we will give an asymptotic formula for $N(\lambda)$.

Theorem 3

$$N(\lambda) = \sum_n \frac{1}{e^{\beta(\lambda_n-\lambda)}+1}, \quad (\beta\to\infty). \tag{11}$$

Proof. Observing that

$$\lim_{\beta\to\infty}\frac{1}{e^{\beta(\lambda_n-\lambda)}+1} = \begin{cases} 1, & \text{when } \lambda_n < \lambda, \\ 0, & \text{when } \lambda_n > \lambda, \end{cases}$$

we have

$$\lim_{\beta\to\infty}\sum_n \frac{1}{e^{\beta(\lambda_n-\lambda)}+1} = \sum_{\lambda_n<\lambda} 1 = N(\lambda).$$

■

Remark 4 *The asymptotic formula for $N(\lambda)$ given by Eq. (11) converts a partial sum ($\sum_{\lambda_n<\lambda}$) into a sum over all possible values ($\sum_{\lambda_n<\infty}$). This will make the calculation somewhat easy.*

In some cases the counting function approximately equals the heat kernel.

Theorem 5 *Let $\rho(\lambda)$ be the number of eigenstates per unit interval (the density of eigenstates). In the limit $\lambda \to \infty$ or $t \to 0$,*

$$N(\lambda) = K\left(\frac{1}{\lambda}\right) \quad or \quad N\left(\frac{1}{t}\right) = K(t), \tag{12}$$

when $\rho(\lambda)$ is a constant.

Proof. In the limit $\lambda \to \infty$ or $t \to 0$, the summations can be converted into integrals:

$$N(\lambda) = \sum_{\lambda_n \leq \lambda} 1 = \int_0^{\lambda} \rho(\lambda') \, d\lambda',$$
$$K(t) = \sum_n e^{-\lambda_n t} = \int_0^{\infty} \rho(\lambda') \, e^{-\lambda' t} d\lambda'.$$

If $\rho(\lambda) = C$, where C is a constant, then

$$N(\lambda) = C\lambda,$$
$$K(t) = \frac{C}{t}.$$

This proves the theorem. ■

This is just the case that Weyl [7], Pleijel [8], and Kac [6] discussed.

SCHOOL OF SCIENCE, TIANJIN UNIVERSITY, TIANJIN, P. R. CHINA
LIUHUI CENTER FOR APPLIED MATHEMATICS, NANKAI UNIVERSITY & TIANJIN UNIVERSITY, TIANJIN, P. R. CHINA
E-*mail address*: daiwusheng@tju.edu.cn

SCHOOL OF SCIENCE, TIANJIN UNIVERSITY, TIANJIN, P. R. CHINA
LIUHUI CENTER FOR APPLIED MATHEMATICS, NANKAI UNIVERSITY & TIANJIN UNIVERSITY, TIANJIN, P. R. CHINA
E-*mail address*: xiemi@tju.edu.cn

References

[1] P. H. Bérard, *Spectral Geometry: Direct and Inverse Problems, With an Appendix by G. Besson*, Springer-Verlag, Berlin, 1986.

[2] M. Berger, *A Panoramic View of Riemannian Geometry*, Springer-Verlag, Berlin, 2003.

[3] J. W. Milnor, Eigenvalues of the Laplace operator on certain manifolds, *Proc. Nat. Acad. Sci. U.S.A.* **51** (1964), 542.

[4] G.Gordon, D. Webb, and S. Wolpert, Isospectral plane domains and surfaces via Riemannian orbifolds, *Invent. Math.* **110** (1992), 1.

[5] V. Ivrii, *Microlocal analysis and precise spectral asymptotics*, Springer-Verlag, Berlin, 1998.

[6] M. Kac, Can one hear the shape of a drum? *Am. Math. Monthly* **73** (1966), 1.

[7] H. Weyl, in: K. Chandrasekharan (Ed.), *Gesammelte Abhandlungen*, Vol. I, Springer, Berlin, 1968.

[8] A. Pleijel, *Ark. Mat.* **2** (1954), 553.